



Sharing Poll Results


Attendees are now viewing the poll results

1. How do you assess ventilator-associated pneumonia in your NICU?

Stop Share Results

Re-launch Polling





Windows-style title bar: Polls [minimize] [maximize] [close]


Sharing Poll Results

Attendees are now viewing the poll results

1. What is the routine in your unit with regards to Lumbar puncture, assuming the infant is stable enough to tolerate the procedure? (Check all that apply) (Multiple choice)

In all infants at the time of performing early-onset sepsis evaluations	(12/100) 12%
<div style="background-color: #0070c0; width: 12%; height: 10px; margin-bottom: 5px;"></div> <hr style="border: 0; border-top: 1px solid #ccc;"/>	
ONLY in infants with positive blood culture during early-onset sepsis evaluations	(60/100) 60%
<div style="background-color: #e69d00; width: 60%; height: 10px; margin-bottom: 5px;"></div> <hr style="border: 0; border-top: 1px solid #ccc;"/>	
In all infants at the time of performing late-onset sepsis evaluations	(45/100) 45%
<div style="background-color: #0070c0; width: 45%; height: 10px; margin-bottom: 5px;"></div> <hr style="border: 0; border-top: 1px solid #ccc;"/>	
ONLY in infants with positive blood culture during late-onset sepsis evaluations (including coagulase-negative Staphylococcus)	(27/100) 27%
<div style="background-color: #0070c0; width: 27%; height: 10px; margin-bottom: 5px;"></div> <hr style="border: 0; border-top: 1px solid #ccc;"/>	
ONLY in infants with positive blood culture during late-onset sepsis evaluations (excluding coagulase-negative Staphylococcus)	(35/100) 35%
<div style="background-color: #0070c0; width: 35%; height: 10px; margin-bottom: 5px;"></div> <hr style="border: 0; border-top: 1px solid #ccc;"/>	

Stop Share ResultsRe-launch Polling



Sharing Poll Results

Attendees are now viewing the poll results

1. Do you use arterial catheter to collect blood culture(s) in infants with suspected central catheter associated blood stream infections (CLABSI) if the baby has an arterial catheter?

Yes (47) 44%



No (60) 56%



Stop Share Results

Re-launch Polling