


Sharing Poll Results


Attendees are now viewing the poll results

1. In my NICU, I would usually provide medical treatment for a hemodynamically significant PDA in preterm infants <29 weeks' GA:



Stop Share Results

Re-launch Polling



Windows title bar: Polls


Sharing Poll Results


Attendees are now viewing the poll results

1. Does your "Golden Hour management" include initiation of TPN and Caffeine?

Yes, all 2 are initiated within Golden hour	(57) 45%
<div style="border: 1px solid #ccc; background-color: #eee; padding: 2px;"><div style="background-color: #e67e22; width: 45%;"></div></div>	
No, these are initiated outside the Golden hour	(15) 12%
<div style="border: 1px solid #ccc; background-color: #eee; padding: 2px;"><div style="background-color: #3498db; width: 12%;"></div></div>	
Some of these are initiated in Golden hour	(56) 44%
<div style="border: 1px solid #ccc; background-color: #eee; padding: 2px;"><div style="background-color: #3498db; width: 44%;"></div></div>	

Stop Share ResultsRe-launch Polling





Windows title bar: Polls

Sharing Poll Results

Attendees are now viewing the poll results

1. Approximately what proportion of infants born AFTER 35 weeks of gestation were exposed to antenatal corticosteroids during the pregnancy?

Response	Count	Percentage
20%	13	10%
30%	2	2%
40%	2	2%
50%	2	2%
60%	0	0%
I do not know	106	85%

[Stop Share Results](#) [Re-launch Polling](#)

Sharing Poll Results

Attendees are now viewing the poll results

1. What is your feeding practice during blood transfusion in extreme preterm infants ? (Pick one)



Stop Share Results

Re-launch Polling